| Audit Review Table Presbyterian Health Plan, Inc. (Org ID: 579, Sub ID: 4104, Medicaid, Spec Area: None, Spec Proj: None, Contract Number: None) Measurement Year - 2022; Date & Timestamp - 6/7/2023 10:06:15 AM | | | | | | |
|---|-----------------|------------------|--------|-------------------|----------------------|--|
| | | | | | | |
| Measure/Data Element Effectiveness of Care | Benefit Offered | Rate | Status | Audit Designation | Comment | |
| Weight Assessment and Counseling for Nutrition and Physical | | | | | | |
| Activity for Children/Adolescents (WCC) | | | | | | |
| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents - BMI percentile (Total) | | 75.31% | R | R | Reported | |
| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents - Counseling for Nutrition (Total) | | 61.60% | R | R | Reported | |
| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents - Counseling for Physical Activity (Total) | | 58.35% | R | R | Reported | |
| Childhood Immunization Status (CIS) | | | | | | |
| Childhood Immunization Status - DTaP | | 75.67% | R | R | Reported | |
| Childhood Immunization Status - IPV | | 87.35% | R | R | Reported | |
| Childhood Immunization Status - MMR | | 89.05% | R | R | Reported | |
| Childhood Immunization Status - HiB | | 85.89% | R | R R | Reported | |
| Childhood Immunization Status - Hepatitis B Childhood Immunization Status - VZV | | 91.73% | R | R | Reported | |
| Childhood Immunization Status - V2V Childhood Immunization Status - Pneumococcal Conjugate | | 85.89% 76.64% | R | R | Reported Reported | |
| Childhood Immunization Status - Friedmococcal Conjugate | | 85.89% | R | R | Reported | |
| Childhood Immunization Status - Repairts A | | 75.91% | R | R | Reported | |
| Childhood Immunization Status - Influenza | | 50.12% | R | R | Reported | |
| Childhood Immunization Status - Combo 3 | | 70.56% | R | R | Reported | |
| Childhood Immunization Status - Combo 7 | | 65.94% | R | R | Reported | |
| Childhood Immunization Status - Combo 10 | | 42.58% | R | R | Reported | |
| Immunizations for Adolescents (IMA) | | | | | | |
| Immunizations for Adolescents - Meningococcal | | 84.18% | R | R | Reported | |
| Immunizations for Adolescents - Tdap | | 86.13% | R | R | Reported | |
| Immunizations for Adolescents - HPV | | 35.77% | R | R | Reported | |
| Immunizations for Adolescents - Combination 1 | | 83.94% 35.52% | R | R R | Reported | |
| Immunizations for Adolescents - Combination 2 Lead Screening in Children (LSC) | | 35.52% | ĸ | ĸ | Reported | |
| Lead Screening in Children (LSC) | | 34.79% | R | R | Reported | |
| Breast Cancer Screening (BCS) | | 04.1070 | IX. | IX. | rteporteu | |
| Breast Cancer Screening | | 39.92% | R | R | Reported | |
| Cervical Cancer Screening (CCS) | | | | | | |
| Cervical Cancer Screening | | 48.66% | R | R | Reported | |
| Colorectal Cancer Screening (COL) | | | | | | |
| Colorectal Cancer Screening (Total) | | 29.64% | R | R | Reported | |
| Chlamydia Screening in Women (CHL) | | | | | | |
| Chlamydia Screening in Women (Total) | | 47.76% | R | R | Reported | |
| Appropriate Testing for Pharyngitis (CWP) | Y | 70 540/ | P | P | Demosteral | |
| Appropriate Testing for Pharyngitis (Total) Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR) | | 76.51% | R | R | Reported | |
| Use of Spirometry Testing in the Assessment and Diagnosis of COPD | | 17.88% | R | R | Reported | |
| Pharmacotherapy Management of COPD Exacerbation (PCE) | Y | | | | | |
| Pharmacotherapy Management of COPD Exacerbation - Systemic Corticosteroid | | 54.42% | R | R | Reported | |
| Pharmacotherapy Management of COPD Exacerbation - Bronchodilator | | 63.67% | R | R | Reported | |
| Asthma Medication Ratio (AMR) | Y | | | | | |
| Asthma Medication Ratio (Total) | | 73.13% | R | R | Reported | |
| Controlling High Blood Pressure (CBP) | | F0 F0% | P | P | Demosteral | |
| Controlling High Blood Pressure Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) | Y | 52.59% | R | R | Reported | |
| Persistence of Beta-Blocker Treatment After a Heart Attack | | 70.06% | R | R | Reported | |
| Statin Therapy for Patients With Cardiovascular Disease (SPC) | Y | | | | | |
| Statin Therapy for Patients With Cardiovascular Disease - Received Statin Therapy (Total) | | 63.27% | R | R | Reported | |
| Statin Therapy (10tal) Statin Therapy for Patients With Cardiovascular Disease - Statin | | 64.770/ | | D | Dam-it-1 | |
| Adherence 80% (Total) | | 64.77% | R | R | Reported | |
| Cardiac Rehabilitation (CRE) | | | | - | _ | |
| Cardiac Rehabilitation - Initiation (Total) | | 0.92% | R | R | Reported | |
| Cardiac Rehabilitation - Engagement1 (Total) | | 3.49% | R | R | Reported | |
| Cardiac Rehabilitation - Engagement2 (Total) Cardiac Rehabilitation - Achievement (Total) | | 3.13% 1.10% | R | R R | Reported Reported | |
| | | 1.1070 | IX IX | 13 | reported | |

| Hemoglobin A1c Control for Patients With Diabetes - HbA1c Control | | | | Γ | |
|--|---|----------------|---|-------|----------------------|
| (<8%) | | 40.88% | R | R | Reported |
| Hemoglobin A1c Control for Patients With Diabetes - Poor HbA1c Control | | 52.31% | R | R | Reported |
| Blood Pressure Control for Patients With Diabetes (BPD) Blood Pressure Control for Patients With Diabetes | | 54.26% | R | R | Reported |
| Eye Exam for Patients With Diabetes (EED) | | 34.2070 | K | IN IN | Reported |
| Eye Exam for Patients With Diabetes | | 39.17% | R | R | Reported |
| Kidney Health Evaluation for Patients With Diabetes (KED) | | | | | |
| Kidney Health Evaluation for Patients With Diabetes (Total) Statin Therapy for Patients With Diabetes (SPD) | Y | 33.51% | R | R | Reported |
| Statin Therapy for Patients With Diabetes - Received Statin Therapy | | 50.54% | R | R | Reported |
| | | | | | |
| Statin Therapy for Patients With Diabetes - Statin Adherence 80% | | 61.35% | R | R | Reported |
| Diagnosed Mental Health Disorders (DMH) Diagnosed Mental Health Disorders (Total) | | 26.34% | R | R | Reported |
| Antidepressant Medication Management (AMM) | Y | 20.3470 | K | | Reported |
| Antidepressant Medication Management - Effective Acute Phase | | 61.99% | R | R | Reported |
| Treatment Antidepressant Medication Management - Effective Continuation | | 40.00% | R | R | |
| Phase Treatment | | 42.33% | R | R | Reported |
| Follow-Up Care for Children Prescribed ADHD Medication (ADD) | Y | | | | |
| Follow-Up Care for Children Prescribed ADHD Medication - Initiation Phase | | 28.73% | R | R | Reported |
| Follow-Up Care for Children Prescribed ADHD Medication - | | 36.73% | R | R | Reported |
| Continuation and Maintenance Phase Follow-Up After Hospitalization for Mental Illness (FUH) | Y | | | | |
| Follow-Up After Hospitalization For Mental Illness - 30 days (Total) | | 55.03% | R | R | Reported |
| Follow-Up After Hospitalization For Mental Illness - 7 days (Total) | | 35.14% | R | R | Reported |
| Follow-Up After Emergency Department Visit for Mental Illness | Y | | | · · | |
| (FUM) Follow-Up After Emergency Department Visit for Mental Illness - 30 | T | | _ | _ | |
| days (Total) Follow-Up After Emergency Department Visit for Mental Illness - 7 | | 59.36% | R | R | Reported |
| days (Total) | | 45.59% | R | R | Reported |
| Diagnosed Substance Use Disorders (DSU) | | | _ | | |
| Diagnosed Substance Use Disorders - Alcohol (Total) Diagnosed Substance Use Disorders - Opioid (Total) | | 4.25% 3.33% | R | R | Reported Reported |
| Diagnosed Substance Use Disorders - Other (Total) | | 4.26% | R | R | Reported |
| Diagnosed Substance Use Disorders - Any (Total) | | 9.14% | R | R | Reported |
| Follow-Up After High-Intensity Care for Substance Use Disorder | Y | | | | |
| (FUI) Follow-Up After High-Intensity Care for Substance Use Disorder - 30 | | 42.04% | R | R | Reported |
| days (Total) Follow-Up After High-Intensity Care for Substance Use Disorder - 7 | | 20.72% | R | R | |
| Days (Total) Follow-Up After Emergency Department Visit for Substance Use | | 20.72% | ĸ | ĸ | Reported |
| (FUA) | Y | | | | |
| Follow-Up After Emergency Department Visit for Substance Use - 30 days (Total) | | 34.98% | R | R | Reported |
| Follow-Up After Emergency Department Visit for Substance Use - 7 | | 23.57% | R | R | Reported |
| days (Total) Pharmacotherapy for Opioid Use Disorder (POD) | Y | | | | |
| Pharmacotherapy for Opioid Use Disorder (Total) | | 17.72% | R | R | Reported |
| Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) | Y | | | | |
| Diabetes Screening for People With Schizophrenia or Bipolar | | 80.59% | R | R | Reported |
| Disorder Who Are Using Antipsychotic Medications Diabetes Monitoring for People With Diabetes and | | | | | |
| Schizophrenia (SMD) | | 64.08% | R | R | Reported |
| Diabetes Monitoring for People With Diabetes and Schizophrenia Cardiovascular Monitoring for People With Cardiovascular | | 64.08% | ĸ | ĸ | Reported |
| Disease and Schizophrenia (SMC) | | | | | |
| Cardiovascular Monitoring for People With Cardiovascular Disease and Schizophrenia | | 73.53% | R | R | Reported |
| Adherence to Antipsychotic Medications for Individuals With | Y | | | | |
| Schizophrenia (SAA) Adherence to Antipsychotic Medications for Individuals With | | FF 00% | P | P | Demonteria |
| Schizophrenia Metabolic Monitoring for Children and Adolescents on | | 55.02% | R | R | Reported |
| Antipsychotics (APM) | Y | | | | |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - Blood Glucose Testing (Total) | | 60.65% | R | R | Reported |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - | | 39.23% | R | R | Reported |
| Cholesterol Testing (Total) Metabolic Monitoring for Children and Adolescents on Antipsychotics - | | 38.19% | R | R | Reported |
| Blood Glucose and Cholesterol Testing (Total) Non-Recommended Cervical Cancer Screening in Adolescent | | | | | |
| Females (NCS) | | | | | |

| Non-Recommended Cervical Cancer Screening in Adolescent Females | | 0.37% | R | R | Reported |
|---|---|----------------------------|-------------|-------------|----------------------|
| Appropriate Treatment for Upper Respiratory Infection (URI) | Y | | | | |
| Appropriate Treatment for Upper Respiratory Infection (Total) | | 84.39% | R | R | Reported |
| Avoidance of Antibiotic Treatment for Acute | Y | | | | |
| Bronchitis/Bronchiolitis (AAB) Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis | • | | | | |
| (Total) | | 58.11% | R | R | Reported |
| Use of Imaging Studies for Low Back Pain (LBP) | | | | | |
| Use of Imaging Studies for Low Back Pain (Total) | | 73.58% | R | R | Reported |
| Use of Opioids at High Dosage (HDO) | Y | | | | |
| Use of Opioids at High Dosage | | 4.12% | R | R | Reported |
| Use of Opioids From Multiple Providers (UOP) | Y | 45.00% | | | Denoted |
| Use of Opioids From Multiple Providers - Multiple Prescribers Use of Opioids From Multiple Providers - Multiple Pharmacies | | 15.09% 1.85% | R R | R | Reported |
| Use of Opioids From Multiple Providers - Multiple Praimacles | | | | | Reported |
| Multiple Pharmacies | | 0.77% | R | R | Reported |
| Risk of Continued Opioid Use (COU) | Y | | | | |
| Risk of Continued Opioid Use - >=15 Days (Total) | | 4.85% | R | R | Reported |
| Risk of Continued Opioid Use - >=31 Days (Total) | | 3.48% | R | R | Reported |
| Access/Availability of Care | | - | | | - |
| Adults' Access to Preventive/Ambulatory Health Services (AAP) | | | | | |
| Adults' Access to Preventive/Ambulatory Health Services (Total) | | 73.39% | R | R | Reported |
| , | Y | | | | . toportod |
| Annual Dental Visit (ADV) Annual Dental Visit (Total) | T | 61.06% | R | R | Reported |
| Initiation and Engagement of Substance Use Disorder Treatment | | 01.00% | 17 | n. | Reported |
| (IET) | Y | | | | |
| Initiation and Engagement of Substance Use Disorder Treatment - Initiation of SUD Treatment - Total (Total) | | 55.10% | R | R | Reported |
| Initiation of SOD Treatment - Total (Total) Initiation and Engagement of Substance Use Disorder Treatment - | | 00.700/ | | P | Demonstructure of |
| Engagement of SUD Treatment - Total (Total) | | 22.73% | R | R | Reported |
| Prenatal and Postpartum Care (PPC) | | | | | |
| Prenatal and Postpartum Care - Timeliness of Prenatal Care | | 78.14% | R | R | Reported |
| Prenatal and Postpartum Care - Postpartum Care Use of First-Line Psychosocial Care for Children and | | 72.95% | R | R | Reported |
| Adolescents on Antipsychotics (APP) | Y | | | | |
| Use of First-Line Psychosocial Care for Children and Adolescents on | | 62.36% | R | R | Reported |
| Antipsychotics (Total) Utilization and Risk Adjusted Utilization | | | | | · · |
| Well-Child Visits in the First 30 Months of Life (W30) | | | | | |
| Well-Child Visits in the First 30 Months of Life (First 15 Months) | | 65.98% | R | R | Reported |
| Well-Child Visits in the First 30 Months of Life (15 Months-30 | | 68.68% | R | R | Reported |
| Months | | 00.0078 | ĸ | IX. | Reported |
| Child and Adolescent Well-Care Visits (WCV) | | 44.450/ | | | Denoted |
| Child and Adolescent Well-Care Visits (Total) | | 44.45% | R R | R | Reported |
| Frequency of Selected Procedures (FSP) Ambulatory Care (AMB) | | | R | R | Reported Reported |
| Inpatient Utilization - General Hospital/Acute Care (IPU) | | | R | R | Reported |
| Antibiotic Utilization for Respiratory Conditions (AXR) | Y | | , N | i k | rioponed |
| Antibiotic Utilization for Respiratory Conditions (Total) | - | 20.61% | R | R | Reported |
| Plan All-Cause Readmissions (PCR) | | | R | R | Reported |
| Health Plan Descriptive Information | | | | | |
| Enrollment by Product Line (ENP) | | | R | R | Reported |
| Language Diversity of Membership (LDM) | | | R | R | Reported |
| Race/Ethnicity Diversity of Membership (RDM) | | | R | R | Reported |
| Measures Reported Using Electronic Clinical Data Systems | | | | | |
| Childhood Immunization Status (CIS-E) | | | _ | - | P |
| Childhood Immunization Status - DTaP | | 52.58% | R | R | Reported |
| Childhood Immunization Status - IPV Childhood Immunization Status - MMR | | 67.80% 72.51% | R R | R | Reported Reported |
| Childhood Immunization Status - MiR Childhood Immunization Status - HiB | | 67.84% | R | R | Reported |
| Childhood Immunization Status - Hepatitis B | | 58.89% | R | R | Reported |
| Childhood Immunization Status - VZV Childhood Immunization Status - VZV | | 71.59% | R | R | Reported |
| Childhood Immunization Status - Pneumococcal Conjugate | | 55.19% | R | R | Reported |
| Childhood Immunization Status - Hepatitis A | | 73.30% | R | R | Reported |
| Childhood Immunization Status - Rotavirus | | 59.29% | R | R | Reported |
| Childhood Immunization Status - Influenza | | 36.07% | R | R | Reported |
| Childhood Immunization Status - Combo 3 | | 40.23% | R | R | Reported |
| Childhood Immunization Status - Combo 7 | | 37.11% | R | R | Reported |
| Childhood Immunization Status - Combo 10 | | 20.84% | R | R | Reported |
| Immunizations for Adolescents (IMA-E) | | | | - | |
| Immunizations for Adolescents - Meningococcal | | 63.92% | R | R | Reported |
| | | | | | |
| Immunizations for Adolescents - Tdap | | 67.36% | R | R | Reported |
| | | 67.36% 26.48% 62.28% | R R R | R R R | Reported Reported |

| Immunizations for Adolescents - Combination 2 | | 25.47% | R | R | Reported |
|---|---|---------|-----|-----|-----------|
| Breast Cancer Screening (BCS-E) | | | | | |
| Breast Cancer Screening | | 38.71% | R | R | Reported |
| Colorectal Cancer Screening (COL-E) | | 00.1170 | N N | - R | rtoportou |
| Colorectal Cancer Screening (COL-L) | | 25.99% | R | R | Reported |
| Follow-Up Care for Children Prescribed ADHD Medication (ADD- | | 25.99% | ĸ | R | Reported |
| E) | Y | | | | |
| Follow-Up Care for Children Prescribed ADHD Medication - Initiation Phase | | 28.73% | R | R | Reported |
| Follow-Up Care for Children Prescribed ADHD Medication - | | 36.73% | R | R | Reported |
| Continuation and Maintenance Phase | | | | | |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM-E) | Y | | | | |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics | | E7.04% | R | P | Demosterd |
| Blood Glucose Testing (Total) | | 57.84% | R | R | Reported |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - | | 31.48% | R | R | Reported |
| Cholesterol Testing (Total) | | | | | |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - Blood Glucose and Cholesterol Testing (Total) | | 30.69% | R | R | Reported |
| Depression Screening and Follow-Up for Adolescents and | | | | | |
| Adults (DSF-E) | | | | | |
| Depression Screening and Follow-Up for Adolescents and Adults - | | 19.72% | R | R | Reported |
| Depression Screening (Total) Depression Screening and Follow-Up for Adolescents and Adults - | | | | | |
| Follow-Up on Positive Screen (Total) | | 74.48% | R | R | Reported |
| Utilization of the PHQ-9 to Monitor Depression Symptoms for | | | | | |
| Adolescents and Adults (DMS-E) | | | | | |
| Utilization of the PHQ-9 to Monitor Depression Symptoms for | | 16.58% | R | R | Reported |
| Adolescents and Adults - Utilization of PHQ-9-Total (Total) Depression Remission or Response for Adolescents and Adults | | | | | |
| (DRR-E) | | | | | |
| Depression Remission or Response for Adolescents and Adults - Follow-Up PHQ-9 (Total) | | 54.87% | R | R | Reported |
| Depression Remission or Response for Adolescents and Adults - Depression Remission (Total) | | 18.58% | R | R | Reported |
| Depression Remission or Response for Adolescents and Adults - | | 26.42% | R | R | Reported |
| Depression Response (Total) | | | | | |
| Unhealthy Alcohol Use Screening and Follow-Up (ASF-E) | | | | | |
| Unhealthy Alcohol Use Screening and Follow-Up - Unhealthy Alcohol Use Screening (Total) | | 0.00% | R | R | Reported |
| Unhealthy Alcohol Use Screening and Follow-Up - Alcohol | | | NA | R | Reported |
| Counseling or Other Follow-Up Care (Total) Adult Immunization Status (AIS-E) | | | | | |
| · · · · · · · · · · · · · · · · · · · | | 9,18% | R | R | Demonteri |
| Adult Immunization Status - Influenza (19-65) | | | | | Reported |
| Adult Immunization Status - Td/Tdap (19-65) | | 32.77% | R | R | Reported |
| Adult Immunization Status - Zoster (50-65) | | 5.47% | R | R | Reported |
| Prenatal Immunization Status (PRS-E) | | | _ | _ | |
| Prenatal Immunization Status - Influenza | | 28.58% | R | R | Reported |
| Prenatal Immunization Status - Tdap | | 57.85% | R | R | Reported |
| Prenatal Immunization Status - Combination | | 24.58% | R | R | Reported |
| Prenatal Depression Screening and Follow-Up (PND-E) | | | | | |
| Prenatal Depression Screening and Follow-Up - Depression Screening | | 36.20% | R | R | Reported |
| Prenatal Depression Screening and Follow-Up - Follow-Up on Positive Screen | | 54.90% | R | R | Reported |
| Postpartum Depression Screening and Follow-Up (PDS-E) | | | | | |
| Postpartum Depression Screening and Follow-Up - Depression Screening | | 19.22% | R | R | Reported |
| Postpartum Depression Screening and Follow-Up - Follow-Up on Positive Screen | | 82.35% | R | R | Reported |