Audit Review Table

Presbyterian Health Plan, Inc. (Org ID: 579, SubID: 4104, Medicaid, Spec Area: None, Spec Proj: None); Measurement Year - 2012

The Auditor lock has been applied to this submission.

Pharmacy Reversal - Pharmacy Reversals Included

Measure/Data Element	Report Measure	Benefit Offered	Rotated Measure	Rate	Reportable	Comment
Effectiveness of Care: Prevention and Screening						
Adult BMI Assessment (aba)	Υ		N	74.70%	R	Reportable
weight Assessment and Counseling for Nutrition and Physical Activity for	Υ		N			
Children/Adolescents (wcc)				00.470/		5
BMI Percentile				39.17%	R	Reportable
Counseling for Nutrition				44.53%	R	Reportable
Counseling for Physical Activity	.,			34.55%	R	Reportable
Childhood Immunization Status (cis)	Y		N		_	
DTaP				85.40%	R	Reportable
IPV				95.86%	R	Reportable
MMR				92.94%	R	Reportable
HiB				96.35%	R	Reportable
Hepatitis B				91.24%	R	Reportable
VZV				92.21%	R	Reportable
Pneumococcal Conjugate				82.97%	R	Reportable
Hepatitis A				85.89%	R	Reportable
Rotavirus				78.10%	R	Reportable
Influenza				58.39%	R	Reportable
Combination #2				80.29%	R	Reportable
Combination #3				74.70%	R	Reportable
Combination #4				71.29%	R	Reportable
Combination #5				64.48%	R	Reportable
Combination #6				50.85%	R	Reportable
Combination #7				62.29%	R	Reportable
Combination #8				48.91%	R	Reportable
Combination #9				45.01%	R	Reportable
Combination #10				43.80%	R	Reportable
Immunizations for Adolescents (ima)	Υ					

Meningococcal				56.93%	R	Reportable
Tdap/Td				72.99%	R	Reportable
Combination #1				54.50%	R	Reportable
Human Papillomavirus Vaccine for Female Adolescents (hpv)	Υ			18.49%	R	Reportable
Lead Screening in Children (Isc)	Υ		N	34.06%	R	Reportable
Breast Cancer Screening (bcs)	Υ			49.39%	R	Reportable
Cervical Cancer Screening (ccs)	Y			60.34%	R	Reportable
Chlamydia Screening in Women (chl)	Υ					
16-20 Years				41.83%	R	Reportable
21-24 Years				61.86%	R	Reportable
Total				47.37%	R	Reportable
Effectiveness of Care: Respiratory Conditions						
Appropriate Testing for Children with Pharyngitis (cwp)	Υ	Y		68.52%	R	Reportable
Appropriate Treatment for Children With URI (uri)	Υ	Y		85.75%	R	Reportable
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (aab)	Υ	Υ		15.99%	R	Reportable
Use of Spirometry Testing in the Assessment and Diagnosis of COPD (spr)	Y			22.00%	R	Reportable
Pharmacotherapy Management of COPD Exacerbation (pce)	Υ	Y				
Systemic Corticosteroid				66.67%	R	Reportable
Bronchodilator				83.87%	R	Reportable
Use of Appropriate Medications for People With Asthma (asm)	Υ	Y				
5-11 Years				82.72%	R	Reportable
12-18 Years				74.68%	R	Reportable
19-50 Years				59.87%	R	Reportable
51-64 Years				59.42%	R	Reportable
Total				76.88%	R	Reportable
Medication Management for People With Asthma (mma)	Υ	Y				

5-11 Years - Medication Compliance 50%				44.27%	R	Reportable
5-11 Years - Medication Compliance 75%				22.18%	R	Reportable
12-18 Years - Medication Compliance 50%				46.01%	R	Reportable
12-18 Years - Medication Compliance 75%				23.94%	R	Reportable
19-50 Years - Medication Compliance 50%				56.91%	R	Reportable
19-50 Years - Medication Compliance 75%				38.83%	R	Reportable
51-64 Years - Medication Compliance 50%				73.17%	R	Reportable
51-64 Years - Medication Compliance 75%				46.34%	R	Reportable
Total - Medication Compliance 50%				46.47%	R	Reportable
Total - Medication Compliance 75%				24.63%	R	Reportable
Asthma Medication Ratio (amr)	Υ	Υ				
5-11 Years				46.79%	R	Reportable
12-18 Years				42.24%	R	Reportable
19-50 Years				28.12%	R	Reportable
51-64 Years				42.03%	R	Reportable
Total				43.07%	R	Reportable
Effectiveness of Care: Cardiovascular						
Cholesterol Management for Patients With						
Cardiovascular	Υ		N			
•						
Conditions (cmc)						
LDL-C Screening Performed				79.52%	R	Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL)				50.00%	R	Reportable
LDL-C Screening Performed	Y					
LDL-C Screening Performed LDL-C Control (<100 mg/dL)	Y			50.00%	R	Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart	Y	Y		50.00%	R	Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart Attack (pbh)		Y		50.00% 64.23%	R R	Reportable Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart Attack (pbh) Effectiveness of Care: Diabetes	Y	Y		50.00% 64.23%	R R	Reportable Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart Attack (pbh) Effectiveness of Care: Diabetes Comprehensive Diabetes Care (cdc)		Y	N	50.00% 64.23% NA	R R R	Reportable Reportable Denominator fewer than 30
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart Attack (pbh) Effectiveness of Care: Diabetes Comprehensive Diabetes Care (cdc) Hemoglobin A1c (HbA1c) Testing	Y	Y	N	50.00% 64.23% NA 85.77%	R R R	Reportable Reportable Denominator fewer than 30 Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart Attack (pbh) Effectiveness of Care: Diabetes Comprehensive Diabetes Care (cdc) Hemoglobin A1c (HbA1c) Testing HbA1c Poor Control (>9.0%)	Y	Y	N	50.00% 64.23% NA 85.77% 43.25%	R R R	Reportable Reportable Denominator fewer than 30 Reportable Reportable Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart Attack (pbh) Effectiveness of Care: Diabetes Comprehensive Diabetes Care (cdc) Hemoglobin A1c (HbA1c) Testing HbA1c Poor Control (>9.0%) HbA1c Control (<8.0%)	Y	Y	N	50.00% 64.23% NA 85.77% 43.25% 47.99%	R R R R	Reportable Reportable Denominator fewer than 30 Reportable Reportable Reportable Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart Attack (pbh) Effectiveness of Care: Diabetes Comprehensive Diabetes Care (cdc) Hemoglobin A1c (HbA1c) Testing HbA1c Poor Control (>9.0%) HbA1c Control (<8.0%) HbA1c Control (<7.0%)	Y	Y	N	50.00% 64.23% NA NA 85.77% 43.25% 47.99% 33.74%	R R R R R R	Reportable Reportable Denominator fewer than 30 Reportable Reportable Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart Attack (pbh) Effectiveness of Care: Diabetes Comprehensive Diabetes Care (cdc) Hemoglobin A1c (HbA1c) Testing HbA1c Poor Control (>9.0%) HbA1c Control (<8.0%) HbA1c Control (<7.0%) Eye Exam (Retinal) Performed	Y	Y	N	50.00% 64.23% NA 85.77% 43.25% 47.99% 33.74% 47.26%	R R R R R R R	Reportable Reportable Denominator fewer than 30 Reportable Reportable Reportable Reportable
LDL-C Screening Performed LDL-C Control (<100 mg/dL) Controlling High Blood Pressure (cbp) Persistence of Beta-Blocker Treatment After a Heart Attack (pbh) Effectiveness of Care: Diabetes Comprehensive Diabetes Care (cdc) Hemoglobin A1c (HbA1c) Testing HbA1c Poor Control (>9.0%) HbA1c Control (<8.0%) HbA1c Control (<7.0%)	Y	Y	N	50.00% 64.23% NA NA 85.77% 43.25% 47.99% 33.74%	R R R R R R	Reportable Reportable Denominator fewer than 30 Reportable Reportable Reportable Reportable Reportable Reportable

Medical Attention for Nephropathy			75.00%	R	Reportable
Blood Pressure Control (<140/80 mm Hg)			46.90%	R	Reportable
Blood Pressure Control (<140/90 mm Hg)			67.15%	R	Reportable
Effectiveness of Care:					
Musculoskeletal					
Disease Modifying Anti-Rheumatic Drug Therapy in Rheumatoid Arthritis (art)	Υ	Y	76.25%	R	Reportable
Use of Imaging Studies for Low Back Pain (lbp)	Υ		76.09%	R	Reportable
Effectiveness of Care: Behavioral Health					
Antidepressant Medication Management (amm)	Υ	N			
Effective Acute Phase Treatment			NB	R	Required benefit not offered
Effective Continuation Phase Treatment			NB	R	Required benefit not offered
Follow-Up Care for Children Prescribed ADHD Medication (add)	Υ	Y			
Initiation Phase			36.48%	R	Reportable
Continuation and Maintenance (C&M) Phase			40.80%	R	Reportable
Follow-Up After Hospitalization for Mental Illness (fuh)	Y	N			
30-Day Follow-Up			NB	R	Required benefit not offered
7-Day Follow-Up			NB	R	Required benefit not offered
Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medication (ssd)	Y	Y	84.55%	R	Reportable
Diabetes Monitoring for People With Diabetes and Schizophrenia (smd)	Υ		76.19%	R	Reportable
Cardiovascular Monitoring for People With Cardiovascular Disease and Schizophrenia (smc)	Υ		NA	R	Denominator fewer than 30
Adherence to Antipsychotic Medications for Individuals With Schizophrenia (saa)	Y	Y	32.69%	R	Reportable

Effectiveness of Care: Medication Management		_	_	_	_	
Annual Monitoring for Patients on Persistent Medications (mpm)	Υ	Y				
ACE Inhibitors or ARBs				85.05%	R	Reportable
Digoxin				NA	R	Denominator fewer than 30
Diuretics				86.46%	R	Reportable
Anticonvulsants				68.82%	R	Reportable
Total				83.30%	R	Reportable
Access/Availability of Care						
Adults' Access to Preventive/Ambulatory Health Services (aap)	Υ					
20-44 Years				84.86%	R	Reportable
45-64 Years				86.76%	R	Reportable
65+ Years				86.49%	R	Reportable
Total				85.35%	R	Reportable
Children and Adolescents' Access to Primary Care Practitioners (cap)	Υ					
12-24 Months				97.15%	R	Reportable
25 Months - 6 Years				88.56%	R	Reportable
7-11 Years				91.47%	R	Reportable
12-19 Years				91.19%	R	Reportable
Annual Dental Visit (adv)	Y	Y				
2-3 Years				55.78%	R	Reportable
4-6 Years				75.19%	R	Reportable
7-10 Years				79.65%	R	Reportable
11-14 Years				74.25%	R	Reportable
15-18 Years				64.12%	R	Reportable
19-21 Years				45.51%	R	Reportable
Total				71.17%	R	Reportable
Initiation and Engagement of AOD Dependence Treatment (iet)	Υ	N				
Initiation of AOD Treatment: 13-17 Years				NB	R	Required benefit not offered
Engagement of AOD Treatment: 13-17 Years				NB	R	Required benefit not offered
Initiation of AOD Treatment: 18+ Years				NB	R	Required benefit not offered

Engagement of AOD Treatment: 18+ Years			NB	R	Required benefit not offered
Initiation of AOD Treatment: Total			NB	R	Required benefit not offered
Engagement of AOD Treatment: Total			NB	R	Required benefit not offered
Prenatal and Postpartum Care (ppc)	Υ				
Timeliness of Prenatal Care			80.05%	R	Reportable
Postpartum Care			57.91%	R	Reportable
Call Answer Timeliness (cat)	Υ		89.34%	R	Reportable
Utilization					
Frequency of Ongoing Prenatal Care (fpc)	Υ				
<21 Percent			11.19%	R	Reportable
21-40 Percent			5.60%	R	Reportable
41-60 Percent			6.81%	R	Reportable
61-80 Percent			20.92%	R	Reportable
81+ Percent			55.47%	R	Reportable
Well-Child Visits in the First 15 Months of Life (w15)	Y	N			
0 Visits			1.95%	R	Reportable
1 Visit			1.22%	R	Reportable
2 Visits			2.19%	R	Reportable
3 Visits			5.11%	R	Reportable
4 Visits			7.30%	R	Reportable
5 Visits			18.98%	R	Reportable
6+ Visits			63.26%	R	Reportable
Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (w34)	Υ	N	58.64%	R	Reportable
Adolescent Well-Care Visits (awc)	Υ	N	45.01%	R	Reportable
Frequency of Selected Procedures (fsp)	Υ			R	Reportable
Ambulatory Care: Total (amba)	Υ			R	Reportable
Ambulatory Care: Dual Eligibles (ambb)	Υ			R	Reportable
Ambulatory Care: Disabled (ambc)	Υ			R	Reportable
Ambulatory Care: Other (ambd)	Υ			R	Reportable
Inpatient UtilizationGeneral Hospital/Acute Care: Total (ipua)	Υ			R	Reportable

Inpatient UtilizationGeneral Hospital/Acute Care: Dual Eligibles (ipub)	Υ		R	Reportable
Inpatient UtilizationGeneral Hospital/Acute Care: Disabled (ipuc)	Υ		R	Reportable
Inpatient UtilizationGeneral Hospital/Acute Care: Other (ipud)	Υ		R	Reportable
Identification of Alcohol and Other Drug Services: Total (iada)	Υ	N	R	Required benefit not offered
Identification of Alcohol and Other Drug Services: Dual Eligibles (iadb)	Υ	N	R	Required benefit not offered
Identification of Alcohol and Other Drug Services: Disabled (iadc)	Υ	N	R	Required benefit not offered
Identification of Alcohol and Other Drug Services: Other (iadd)	Υ	N	R	Required benefit not offered
Mental Health Utilization: Total (mpta)	Υ	N	R	Required benefit not offered
Mental Health Utilization: Dual Eligibles (mptb)	Υ	N	R	Required benefit not offered
Mental Health Utilization: Disabled (mptc)	Υ	N	R	Required benefit not offered
Mental Health Utilization: Other (mptd)	Υ	N	R	Required benefit not offered
Antibiotic Utilization: Total (abxa)	Υ	Υ	R	Reportable
Antibiotic Utilization: Dual Eligibles (abxb)	Υ	Y	R	Reportable
Antibiotic Utilization: Disabled (abxc)	Υ	Y	R	Reportable
Antibiotic Utilization: Other (abxd)	Υ	Y	R	Reportable
Relative Resource Use				
Relative Resource Use for People With Diabetes (rdi)	Υ		R	Reportable
Relative Resource Use for People With Asthma (ras)	Υ	Υ	R	Reportable
Relative Resource Use for People With Cardiovascular Conditions (rca)	Υ		R	Reportable
Relative Resource Use for People With Hypertension (rhy)	Υ		R	Reportable
Relative Resource Use for People With COPD (rco)	Υ		R	Reportable

Health Plan Descriptive Information				
Board Certification (bcr)	Υ		R	Reportable
Total Membership (tlm)	Υ		R	Reportable
Enrollment by Product Line: Total (enpa)	Υ		R	Reportable
Enrollment by Product Line: Dual Eligibles (enpb)	Υ		R	Reportable
Enrollment by Product Line: Disabled (enpc)	Y		R	Reportable
Enrollment by Product Line: Other (enpd)	Υ		R	Reportable
Enrollment by State (ebs)	Υ		R	Reportable
Race/Ethnicity Diversity of Membership (rdm)	Υ		R	Reportable
Language Diversity of Membership (Idm)	Y		R	Reportable
Weeks of Pregnancy at Time of Enrollment in MCO (wop)	Υ		R	Reportable